



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

effects at this point, whether at the tip or in the cambium. But as soon as the cells cease to be meristematic, or embryonic, and begin to form vacuoles the haustoria are formed and modification of the cells begins.

The hyphæ keep up with the growth of the shoots by following the trachæ, from which they penetrate the surrounding tissues.

PHYSIOLOGICAL EFFECTS OF BORDEAUX MIXTURE

It has been claimed that Bordeaux mixture, in addition to its fungicidal effects, augments the assimilative activity of plants on which it is sprayed. Ewart (*Zeitschr. Pflanzenkrank.* XXII, p. 257: *Bot. Gaz.*, June, 1913) finds by experimenting with potatoes, radishes and beans that the yield was always decreased by covering the leaves with the mixture, and in proportion to the strength of the mixture. He also found that the sugar content of currants was increased by spraying the *fruit* with the mixture, and decreased by spraying the leaves alone.

HERMAPHRODITISM IN AMPHIOXUS

Goodrich (*Anat. Anz.*, 1913, p. 318) describes an interesting abnormality in this animal. A male specimen with 25 testes on one side had one of the 25 gonads on the other side a perfectly developed ovary with numbers of large ova. All the 49 testes were perfect and full of sperm.

RESISTANCE IN HIBERNATING ANIMALS

Bertarelli (*Centr. Bakt., 1te Abt. Orig.* XVIII, 1913, p. 566) finds that marmots are not more resistant to rabies, anthrax, tetanus, and diphtheria during hibernation than at other times. Blanchard had previously reported these animals to have increased resistance, during hibernation, to cobra venom, diphtheria, tetanus, trypanosomes and trichina.

MICROSCOPIC MEASUREMENT BY CAMERA LUCIDA

Joly (*Sci. Proc. Roy. Dub. Soc.*, XIII, 1913, p. 441) suggests a simple method for measuring microscopic objects by means of the camera-lucida. Draw two fine lines, diverging from a point, on a